

New Physics bounds from the combination of CKM-universality and high energy data*

MARTÍN GONZÁLEZ-ALONSO

Departament de Física Teòrica and IFIC, Universitat de València-CSIC,
Apt. Correus 22085, E-46071 València, Spain

Through an effective field theory approach, we analyze the new physics (NP) corrections to muon and beta decays and their effects on the extractions of V_{ud} and V_{us} . Assuming nearly flavor blind NP interactions, we find that the only quantity sensitive to NP is $\Delta_{\text{CKM}} \equiv |V_{ud}|^2 + |V_{us}|^2 + |V_{ub}|^2 - 1$, that receives contributions from four short distance operators. The phenomenological bound $\Delta_{\text{CKM}} = (-1 \pm 6) \times 10^{-4}$ provides strong constraints on all four operators, corresponding to an effective scale $\Lambda > 11$ TeV (90% CL). Depending on the operator, this constraint is at the same level or better than that generated by the Z pole observables.

1. Introduction

Thanks to the precise experimental measurements [2] and the theoretical improvements [3], semileptonic (SL) decays of light quarks are a deep probe of the nature of weak interactions [4, 5]. In particular, the precise determination of the elements V_{ud} and V_{us} of the Cabibbo-Kobayashi-Maskawa (CKM) [6] matrix enables tests of the CKM unitarity condition¹ $|V_{ud}|^2 + |V_{us}|^2 + |V_{ub}|^2 = 1$, at the level of 0.001 or better. Assuming that NP contributions scale as $\alpha/\pi(M_W^2/\Lambda^2)$, this test probes energy scales Λ on the order of the TeV, which will be directly probed at the LHC.

While the consequences of Cabibbo universality tests have been considered in some explicit Standard Model (SM) extensions [7], our goal is to study it in a model-independent way. We have to analyze the NP contributions to the muon decay (where the G_F is extracted) and to the different channels that are used to extract the product $G_F V_{ud,us}$. Currently the determinations of V_{ud} and V_{us} are dominated by super-allowed nuclear beta

* Presented at the FLAVIANet topical workshop “Low energy constraints on extensions of the Standard Model” (Kazimierz, Poland), July 2009, based on [1].

¹ $V_{ub} \sim 10^{-3}$ contributes negligibly to this relation.

decays [8] ($V_{ud} = 0.97425(22)$) and K_{l3} decays [9] ($V_{us} = 0.2252(9)$) respectively, although experimental and theoretical improvements in other channels can make them competitive in the near future.

2. Weak scale effective lagrangian

In order to analyze in a model-independent framework NP contributions to both electroweak precision observables (EWPO) and beta decays we take the SM (including the Higgs) as the low-energy limit of a more fundamental theory, and more specifically we assume that: (i) there is a gap between the weak scale v and the NP scale Λ where new degrees of freedom appear; (ii) the NP at the weak scale is weakly coupled, so the electroweak (EW) gauge symmetry is linearly realized; (iii) the violation of total lepton and baryon number is suppressed by a scale much higher than $\Lambda \sim \text{TeV}$. These assumptions lead us to an effective non-renormalizable lagrangian of the form [10]:

$$\mathcal{L}^{(\text{eff})} = \mathcal{L}_{\text{SM}} + \frac{1}{\Lambda} \mathcal{L}_5 + \frac{1}{\Lambda^2} \mathcal{L}_6 + \frac{1}{\Lambda^3} \mathcal{L}_7 + \dots \quad (1)$$

where $\mathcal{L}_n = \sum_i \alpha_i^{(n)} \mathcal{O}_i^{(n)}$, being $\mathcal{O}_i^{(n)}$ local gauge-invariant operators of dimension n built out of SM fields. It can be shown that under the above assumptions, there are no corrections to the SM lagrangian at dimension five, whereas seventy-seven operators appear at dimension six [1, 10], where we truncate the expansion. In order to be consistent with this truncation we will work at linear order in the NP corrections.

For the EWPO *and* beta decays it can be shown that we only need a twenty-five operator basis, with twenty-one $U(3)^5$ invariant and four non-invariant² (we will see the usefulness of this separation later). Only nine of those operators will contribute to the beta and muon decays:

$$O_{ll}^{(1)} = \frac{1}{2} (\bar{l} \gamma^\mu l) (\bar{l} \gamma_\mu l) \quad O_{ll}^{(3)} = \frac{1}{2} (\bar{l} \gamma^\mu \sigma^a l) (\bar{l} \gamma_\mu \sigma^a l), \quad (2)$$

$$O_{lq}^{(3)} = (\bar{l} \gamma^\mu \sigma^a l) (\bar{q} \gamma_\mu \sigma^a q), \quad (3)$$

$$O_{\varphi l}^{(3)} = i(h^\dagger D^\mu \sigma^a \varphi) (\bar{l} \gamma_\mu \sigma^a l) + \text{h.c.}, \quad O_{\varphi q}^{(3)} = i(\varphi^\dagger D^\mu \sigma^a \varphi) (\bar{q} \gamma_\mu \sigma^a q) + \text{h.c.}, \quad (4)$$

$$O_{qde} = (\bar{l} e) (\bar{d} q) + \text{h.c.}, \quad (5)$$

$$O_{lq} = (\bar{l}_a e) \epsilon^{ab} (\bar{q}_b u) + \text{h.c.} \quad O_{lq}^t = (\bar{l}_a \sigma^{\mu\nu} e) \epsilon^{ab} (\bar{q}_b \sigma_{\mu\nu} u) + \text{h.c.} \quad (6)$$

$$O_{\varphi\varphi} = i(\varphi^T \epsilon D_\mu \varphi) (\bar{u} \gamma^\mu d) + \text{h.c.}, \quad (7)$$

where only the first five are $U(3)^5$ -invariant.

² We refer to the $U(3)^5$ flavor symmetry of the SM gauge lagrangian (the freedom to make $U(3)$ rotations in family space for each of the five fermionic gauge multiplets).

3. Effective lagrangian for μ and quark β decays

Deriving the low-energy effective lagrangian that describes the muon and beta decays (see ref. [1] for details) we find

$$\mathcal{L}_\mu = \frac{-g^2}{2m_W^2} \left[(1 + \tilde{v}_L) \cdot \bar{e}_L \gamma_\mu \nu_{eL} \bar{\nu}_{\mu L} \gamma^\mu \mu_L + \tilde{s}_R \cdot \bar{e}_R \nu_{eL} \bar{\nu}_{\mu L} \mu_R \right] + h.c. \quad (8)$$

$$\tilde{v}_L = 2 [\hat{\alpha}_{\phi l}^{(3)}]_{11+22^*} - [\hat{\alpha}_{ll}^{(1)}]_{1221} - 2[\hat{\alpha}_{ll}^{(3)}]_{1122-\frac{1}{2}(1221)} \quad (9)$$

$$\tilde{s}_R = +2[\hat{\alpha}_{le}]_{2112} , \quad (10)$$

$$\begin{aligned} \mathcal{L}_{d_j} = \frac{-g^2}{2m_W^2} V_{ij} \left[\left(1 + [v_L]_{\ell\ell ij} \right) \bar{\ell}_L \gamma_\mu \nu_{\ell L} \bar{u}_L^i \gamma^\mu d_L^j + [v_R]_{\ell\ell ij} \bar{\ell}_L \gamma_\mu \nu_{\ell L} \bar{u}_R^i \gamma^\mu d_R^j \right. \\ \left. + [s_L]_{\ell\ell ij} \bar{\ell}_R \nu_{\ell L} \bar{u}_R^i d_L^j + [s_R]_{\ell\ell ij} \bar{\ell}_R \nu_{\ell L} \bar{u}_L^i d_R^j \right. \\ \left. + [t_L]_{\ell\ell ij} \bar{\ell}_R \sigma_{\mu\nu} \nu_{\ell L} \bar{u}_R^i \sigma^{\mu\nu} d_L^j \right] + h.c. , \quad (11) \end{aligned}$$

$$V_{ij} \cdot [v_L]_{\ell\ell ij} = 2 V_{ij} [\hat{\alpha}_{\phi l}^{(3)}]_{\ell\ell} + 2 V_{im} [\hat{\alpha}_{\phi q}^{(3)}]_{jm}^* - 2 V_{im} [\hat{\alpha}_{lq}^{(3)}]_{\ell\ell m j} \quad (12)$$

$$V_{ij} \cdot [v_R]_{\ell\ell ij} = -[\hat{\alpha}_{\phi\phi}]_{ij} \quad (13)$$

$$V_{ij} \cdot [s_L]_{\ell\ell ij} = -[\hat{\alpha}_{lq}]_{\ell\ell ji}^* \quad (14)$$

$$V_{ij} \cdot [s_R]_{\ell\ell ij} = -V_{im} [\hat{\alpha}_{qde}]_{\ell\ell jm}^* \quad (15)$$

$$V_{ij} \cdot [t_L]_{\ell\ell ij} = -[\hat{\alpha}_{lq}^t]_{\ell\ell ji}^* . \quad (16)$$

4. Flavor structure of the effective couplings

So far we have not made any assumption about the flavor structures in the couplings $[\hat{\alpha}_X]_{abcd}$. However flavor changing neutral current (FCNC) processes forbid generic structures if $\Lambda \sim \text{TeV}$ and therefore we organize the discussion in terms of perturbations around the $U(3)^5$ flavor symmetry limit, where no problem arises with FCNC.

In the $U(3)^5$ -limit the expressions greatly simplify and all the NP effects can be encoded into the following redefinitions

$$G_F^\mu = (G_F)^{(0)} \left(1 + 4 \hat{\alpha}_{\phi l}^{(3)} - 2 \hat{\alpha}_{ll}^{(3)} \right) , \quad (17)$$

$$G_F^{\text{SL}} = (G_F)^{(0)} \left(1 + 2 \left(\hat{\alpha}_{\phi l}^{(3)} + \hat{\alpha}_{\phi q}^{(3)} - \hat{\alpha}_{lq}^{(3)} \right) \right) , \quad (18)$$

where $G_F^{(0)} = g^2/(4\sqrt{2}m_W^2)$. Consequently we will have

$$V_{ij}^{(\text{pheno})} = V_{ij} \left[1 + 2 \left(\hat{\alpha}_{ll}^{(3)} - \hat{\alpha}_{lq}^{(3)} - \hat{\alpha}_{\phi l}^{(3)} + \hat{\alpha}_{\phi q}^{(3)} \right) \right] , \quad (19)$$

as phenomenological values of $V_{ud,us}$. This shift is independent of the channel used to extract $V_{ud,us}$ and the only way to expose NP contributions is to construct universality tests, in which the absolute normalization of V_{ij} matters. Therefore the NP effects are entirely captured by the quantity

$$\Delta_{\text{CKM}} \equiv |V_{ud}^{(\text{pheno})}|^2 + |V_{us}^{(\text{pheno})}|^2 + |V_{ub}^{(\text{pheno})}|^2 - 1, \quad (20)$$

that in our framework takes the value

$$\Delta_{\text{CKM}} = 4 \left(\hat{\alpha}_{ll}^{(3)} - \hat{\alpha}_{lq}^{(3)} - \hat{\alpha}_{\varphi l}^{(3)} + \hat{\alpha}_{\varphi q}^{(3)} \right). \quad (21)$$

The Minimal Flavor Violation (MFV) hypothesis requires that $U(3)^5$ symmetry is broken in the underlying model only by structures proportional to the SM Yukawa couplings [11], and structures generating neutrino masses [12]. But in MFV the coefficients parameterizing deviations from the $U(3)^5$ -limit are highly suppressed [1] and so we expect the conclusions of the previous subsection to hold. The elements V_{ij} receive a common dominant shift plus suppressed channel-dependent corrections.

In a more general framework the situation can be different because the channel-dependent shifts to V_{ij} could be appreciable and Δ_{CKM} would depend on the channels used. Work in this direction is in progress.

5. Δ_{CKM} versus precision EW measurements

In the limit of approximate $U(3)^5$ invariance, we have shown that Δ_{CKM} constrains a specific combination of the coefficients, that also contribute to the EWPO [13], together with the remaining seventeen operators that make up the $U(3)^5$ invariant sector of our TeV scale effective lagrangian.

The analysis of Han and Skiba [13], that studied the constraints on the same set of twenty-one $U(3)^5$ invariant operators via a global fit to the EWPO, allows us to compare the bound on Δ_{CKM} that we get from them

$$-9.5 \times 10^{-3} \leq \Delta_{\text{CKM}} \leq 0.1 \times 10^{-3} \quad (90\% \text{ C.L.}) , \quad (22)$$

with the direct experimental bound $|\Delta_{\text{CKM}}| \leq 1. \times 10^{-3}$ (90% C.L.) [9]. We see that EWPO leave room for a sizable non-zero Δ_{CKM} and consequently we have to include the direct Δ_{CKM} constraint in the global fit to improve the bounds on NP-couplings (see results in Fig. 1). We see that the main effect is to strengthen the constraints on $O_{lq}^{(3)}$.

In Fig. 2 we show the bounds if we assume a single operator dominance. For all the CKM-operators the direct Δ_{CKM} measurement provides competitive constraints and in the case of $O_{lq}^{(3)}$ the improvement is remarkable. If a non-zero Δ_{CKM} is observed, in the single-operator framework it would be correlated to deviations from the SM expectation in other observables as well. These correlations have been studied in ref. [1].




Fig. 1. 90% C.L. regions projections, using the high energy observables (HEP), the current Δ_{CKM} constraint or an alternative value of $\Delta_{\text{CKM}} = -0.0025 \pm 0.0006$.




Fig. 2. 90% C.L. regions in the single operator analysis. The first row displays the constraint from EWPO and the second row those coming only from Δ_{CKM} .

6. Conclusions

In a model-independent framework and assuming nearly flavor blind NP interactions, it has been shown that the extraction of $V_{ud,us}$ is channel independent and the only NP probe is Δ_{CKM} , that receives contributions from four short distance operators: $O_{ll,lq,\phi l,\phi q}^{(3)}$.

It has been shown that Cabibbo universality tests provide constraints on NP that currently cannot be obtained from other EW precision tests and

collider measurements. The Δ_{CKM} constraint bounds the effective NP scale of all four CKM-operators to be $\Lambda > 11$ TeV (90 % C.L.), what for $O_{lq}^{(3)}$ is almost one order of magnitude stronger than EWPO-bound. Equivalently, should V_{ud} and V_{us} move from their current central values [5], EWPO data would leave room for sizable deviations from quark-lepton universality.

Acknowledgments

I thank V. Cirigliano and J. Jenkins for their help and the LANL T-2 Group for its hospitality and partial support. Work supported by the EU RTN network FLAVIANet [MRTN-CT-2006-035482] and MICINN, Spain [FPU No. AP20050910, FPA2007-60323 and CSD2007-00042 -CPAN-].

REFERENCES

- [1] V. Cirigliano, M. Gonzalez-Alonso and J. Jenkins, arXiv:0908.1754.
- [2] C. Amsler *et al.* [Particle Data Group], Phys. Lett. B **667** (2008) 1.
- [3] W. J. Marciano and A. Sirlin, Phys. Rev. Lett. **96**, 032002 (2006); V. Cirigliano *et al.*, Eur. Phys. J. C **23**, 121 (2002); V. Cirigliano, H. Neufeld and H. Pichl, Eur. Phys. J. C **35**, 53 (2004); V. Cirigliano, M. Giannotti and H. Neufeld, JHEP **0811**, 006 (2008); J. Bijnens and P. Talavera, Nucl. Phys. B **669**, 341 (2003); M. Jamin, J. A. Oller and A. Pich, JHEP **0402**, 047 (2004); V. Cirigliano *et al.*, JHEP **0504**, 006 (2005); D. Becirevic *et al.*, Nucl. Phys. B **705**, 339 (2005); C. Dawson *et al.*, Phys. Rev. D **74**, 114502 (2006); P. A. Boyle *et al.*, Phys. Rev. Lett. **100**, 141601 (2008); V. Lubicz *et al.*, arXiv:0906.4728.
- [4] W. J. Marciano, PoS **KAON** (2008) 003.
- [5] M. Antonelli *et al.* [FlaviaNet WG on Kaon Decays], arXiv:0801.1817.
- [6] N. Cabibbo, Phys. Rev. Lett. **10** (1963) 531; M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49** (1973) 652.
- [7] R. Barbieri *et al.*, Phys. Lett. B **156**, 348 (1985); W. J. Marciano and A. Sirlin, Phys. Rev. D **35** (1987) 1672; K. Hagiwara, S. Matsumoto and Y. Yamada, Phys. Rev. Lett. **75**, 3605 (1995); A. Kurylov and M. J. Ramsey-Musolf, Phys. Rev. Lett. **88**, 071804 (2002)
- [8] J. C. Hardy and I. S. Towner, Phys. Rev. C **79**, 055502 (2009)
- [9] M. Antonelli *et al.*, arXiv:0907.5386 [hep-ph].
- [10] W. Buchmuller and D. Wyler, Nucl. Phys. B **268** (1986) 621.
- [11] R. S. Chivukula and H. Georgi, Phys. Lett. B **188** (1987) 99; L. J. Hall and L. Randall, Phys. Rev. Lett. **65** (1990) 2939; A. J. Buras *et al.*, Phys. Lett. B **500**, 161 (2001); G. D'Ambrosio *et al.*, Nucl. Phys. B **645**, 155 (2002).
- [12] V. Cirigliano *et al.*, Nucl. Phys. B **728**, 121 (2005)
- [13] Z. Han and W. Skiba, Phys. Rev. D **71**, 075009 (2005)